

Meeting abstract

Open Access

Transcriptional profiling of CD4 T-lymphocytes reveals abnormal gene expression in young prediabetic NOD mice

Dorothy N Kakoola, Nataliya I Lenchik, Anita Curcio-Brint and Ivan C Gerling*

Address: Department of Medicine, University of Tennessee Health Science Center and Research Service, Veterans Affairs Medical Center, Memphis, TN 38104, USA

Email: Ivan C Gerling* - igerling@utmem.edu

* Corresponding author

from UT-ORNL-KBRIN Bioinformatics Summit 2009
Pikeville, TN, USA. 20–22 March 2009

Published: 25 June 2009

BMC Bioinformatics 2009, 10(Suppl 7):A15 doi:10.1186/1471-2105-10-S7-A15

This abstract is available from: <http://www.biomedcentral.com/1471-2105/10/S7/A15>

© 2009 Kakoola et al; licensee BioMed Central Ltd.

Background

CD4 T-lymphocytes play a prominent role in Type 1 DM. The NOD mouse is a model for this autoimmune disease. The earliest histological signs of autoimmunity in female NOD mice occur at approximately 5 weeks of age. We hypothesized that this process may be preceded by molecular events in the CD4 T-lymphocytes. To test this hypothesis, we analyzed the transcriptomes of CD4 T-lymphocytes of young "prepathologic" NOD mice.

Methods

Using a magnetic separation system (Miltenyi Biotec), we isolated CD4 T-lymphocytes from spleen leukocytes of female NOD mice, and two control strains, NOR and C57Bl/6, at 2, 3, and 4 weeks of age ($n = 5$ for each strain, except for NOD 2 wk where $n = 4$). CD4 T-lymphocyte transcriptomes were characterized on Affymetrix Mouse 430_2 expression arrays probing ~39,000 mouse genes/ESTs. Gene lists produced from statistical analyses were mined using Ingenuity Pathway Analysis <http://www.ingenuity.com>.

Results

Analysis by 1-way ANOVA ($p < 0.005$, Benjamin-Hochberg Multiple Test Correction (MTC) defined 311, 624, and 581 probesets (genes) with highly significant expression differences between strains at 2, 3 and 4 weeks, respectively. Hierarchical clustering of these probeset lists

identified 58, 115, and 65 probesets, respectively for 2, 3, and 4 week old mice, as being differentially expressed in NOD mice relative to both control strains (17, 32, and 9 were upregulated in NOD mice while 41, 83, and 56 were downregulated). Pathway analysis of the 3 gene lists (58, 115 and 64 probesets) revealed that 7 (HNF4A, IFNG, IL4, TP53, BCL2L1, IL15, and prostaglandin E2) of the top 15 central genes in each network were common to all age groups. IL6 was central in networks of 2 and 4 week old mice but not of 3 week old mice. Uniquely central to individual networks were: MYC, JUN and HRAS at 2 weeks; TNF, NFKB, p38MAPK, and TGFB1 at 3 weeks; and Interferon alpha, IL12 and STAT4 at 4 weeks.

Conclusion

We conclude that young, prediabetic NOD mice have basic (i.e. common, most likely genetic) molecular defects in CD4 T-cells associated with apoptosis/cellular proliferation, Th1-Th2 differentiation and cytokine signaling. Furthermore, abnormalities in apoptosis/cellular proliferation predominate at 2 weeks of age while those in cellular activation/acute phase response signaling and inflammation predominate in 3 week old mice. Finally, abnormalities in the innate immune response and its link to the adaptive immune response are evident at 4 weeks of age.

Acknowledgements

This work was supported by a research supplement grant from the National Institute of Diabetes & Digestive & Kidney Diseases, DK062103.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

