

Poster presentation

Open Access

## On the origin of systematic errors in a simple navigation task

Stefan Glasauer\*<sup>1,2</sup>, Klaus Jahn<sup>3</sup>, Alexandra Stein<sup>2</sup> and Thomas Brandt<sup>1,2</sup>

Address: <sup>1</sup>Clinical Neurosciences, Ludwig-Maximilian University, Munich, Germany, <sup>2</sup>Bernstein Center of Computational Neuroscience Munich, Germany and <sup>3</sup>Neurology, Ludwig-Maximilian University, Munich, Germany

Email: Stefan Glasauer\* - [sglasauer@lmu.de](mailto:sglasauer@lmu.de)

\* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS\*2009  
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):P13 doi:10.1186/1471-2202-10-S1-P13

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P13>

© 2009 Glasauer et al; licensee BioMed Central Ltd.

One of the simplest tasks in navigation is homing: returning to the point of origin after an excursion. If no positional information is given, e.g., in the dark, path integration is used for homing by various species, including ants and humans, to estimate traveled distance from self-motion cues. In desert ants, a systematic undershoot in homing performance, which increased with outbound distance, was contributed to the inherent leakiness of the ant's path integrator [1]. Previous studies on locomotor homing in blindfolded humans [2,3] similarly showed decreasing homing gain for increasing outbound distance, but the amount of error for a particular distance varied among studies. In contrast to the prediction of leaky integration, the gain-distance relation appeared to depend on the range of distances employed with gains larger than unity for short distances and smaller for long distances.

We first verified that this observation still holds for farther distances not yet tested (up to 250 m). Next, we showed that for a similar path integration task – blindfolded goal-directed locomotion – the gain-distance dependence did not hold as would be expected from a leaky path integrator. After thus having excluded the hypothesis that the gain-distance dependence was caused by leaky integration, we developed a new theory: homing distance is determined by Bayesian inference based on prior expectation and current measurement of outbound distance. Comparison of our homing data with that of the previous studies revealed that the dependence of homing distance on outbound distance follows a classical power law relation [4]: a linear slope in double-log representation.

We show that the power-law dependence is compatible with distance representation and Bayesian inference on an internal logarithmic scale, thus unifying two classical psychophysical theories, the Weber-Fechner law and Steven's law [4], with Bayesian inference. Furthermore, our theory provides a reason for the observed range effect and the "central tendency." Notably, a recent study showed the validity of Weber's law for locomotor path integration [5].

To answer the question of how the prior is determined – from cues such as the size of the experimental room or online during the experiment – we performed yet another experiment: two groups of subjects were tested with the same set of distances, but in reverse order. We found that the performance strongly depended on order of presentation, and that this dependence could be explained by Bayesian estimation of the prior implemented as discrete Kalman filter. In summary, we propose a three-stage mechanism for homing in humans: 1) representation of walked distance is affected by amplitude-dependent noise, 2) homing distance is determined by Bayesian integration of the actual distance estimate and prior expectation and 3) the distance prior is estimated online during the experiment by optimal Bayesian inference.

### Acknowledgements

Study supported by BMBF (BCCN Munich 01GQ0440).

### References

1. Sommer S, Wehner R: **The ant's estimation of distance travelled: experiments with desert ants, *Cataglyphis fortis*.** *J Comp Physiol A* 2004, **190**:1-6.

2. Klatzky RL, Loomis JM, Gooledge RG, Cicinelli JG, Doherty S, Pellegrino JW: **Acquisition of route and survey knowledge in the absence of vision.** *J Mot Behav* 1990, **22**:19-43.
3. Schwartz M: **Haptic perception of the distance walked when blindfolded.** *J Exp Psychol Hum Percept Perform* 1999, **25**:852-865.
4. Stevens SS: **On the psychophysical law.** *Psychol Rev* 1957, **64**:153-181.
5. Durgin FH, Akagi M, Gallistel CR, Haiken W: **The precision of locomotor odometry in humans.** *Exp Brain Res* 2008 in press.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

