

Poster presentation

Open Access

## Interplay between spike-timing-dependent plasticity and neuronal correlations gives rise to network structure

Matthieu Gilson\*<sup>1,2,3</sup>, Doreen A Thomas<sup>1,3</sup>, Anthony N Burkitt<sup>1,2</sup>, David B Grayden<sup>1,2</sup> and J Leo van Hemmen<sup>4</sup>

Address: <sup>1</sup>Department of Electrical and Electronic Engineering, University of Melbourne, Melbourne, VIC 3010, Australia, <sup>2</sup>The Bionic Ear Institute, East Melbourne, VIC 3002, Australia, <sup>3</sup>NICTA, Victoria Research Lab, University of Melbourne, Melbourne, VIC 3010, Australia and <sup>4</sup>Physik Department (T35) and BCCN-Munich, Universität München, 85747 Garching bei München, Germany

Email: Doreen A Thomas - doreen.thomas@unimelb.edu.au

\* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS\*2009  
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, **10**(Suppl 1):P200 doi:10.1186/1471-2202-10-S1-P200

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P200>

© 2009 Gilson et al; licensee BioMed Central Ltd.

### Introduction

Spike-Timing-Dependent Plasticity (STDP) describes the change in the strength (or weight) of neuronal connections depending on the pre- and post-synaptic firing activity [1-3], which in turn changes the spiking properties of the post-synaptic neurons on a slower time scale. The interplay between the parameters of the neurons, the plasticity rule and the input stimulation results in changes to the network structure by strengthening or weakening pathways. This leads to functional changes of the neuronal information processing that takes place in the network. In addition to learning spiking-rate-based information, STDP captures the effect of spike-time correlations on short time scales, down to milliseconds, which are neglected by rate-based learning. STDP has been demonstrated to capture, for example, interaural time differences [1] and the presence of narrowly distributed correlations among input spike trains [2].

### Methods

Extending a previously developed framework [3], we consider the effect of STDP in recurrent neuronal networks stimulated by pools of external inputs, where either the input or recurrent connections (or both) are plastic. We use a theoretical framework based on the Poisson neuron model to analytically describe the network dynamics (firing rates and spike-time correlations) and thus the evolu-

tion of the synaptic weights. This framework incorporates the time course of the post-synaptic potentials and synaptic delays. Our analysis focuses on the asymptotic states of a network stimulated by two homogeneous pools of "steady" inputs, namely Poisson spike trains which have fixed firing rates and spike-time correlations.

### Results and discussion

We show that, in all cases, STDP can generate two simultaneous effects on the weight dynamics: a homeostatic equilibrium that stabilizes neuronal activity (firing rates) and a diverging behavior according to the pairwise spike-time correlation structure. Our analysis suggests that weight-dependent STDP may be necessary in order to obtain stability when both the input and recurrent weights are plastic. At the scale of a neuronal network, inhomogeneities in the connectivity and in the input structure results in an interesting behavior where feed-forward synaptic pathways and areas with reinforced feedback emerge in the initially homogeneous recurrent network. This robust weight competition gives rise to neuronal specialization, which can be linked to self-organization [4] that takes place in the primary visual cortex during the first weeks of new-born mammals [5]. Our results support the idea that spike-time correlations play a major role in the functional organization resulting from synaptic learning in networks.

## Acknowledgements

This work was funded partly by NICTA, The Bionic Ear Institute, the Australian Research Council (ARC Discovery Project #DP0771815) and BCCN-Munich.

## References

1. Gerstner W, Kempter R, van Hemmen JL, Wagner H: **A neuronal learning rule for sub-millisecond temporal coding.** *Nature* 1996, **383**:76-78.
2. Kempter R, Gerstner W, van Hemmen JL: **Hebbian learning and spiking neurons.** *Phys Rev E* 1999, **5**:4498-4514.
3. Burkitt AN, Gilson M, van Hemmen JL: **Spike-timing-dependent plasticity for neurons with recurrent connections.** *Biol Cybern* 2007, **96**:533-546.
4. Kohonen T: **Self-organized formation of topologically correct feature maps.** *Biol Cybern* 1982, **43**:59-69.
5. Hubel DH, Wiesel TN: **Receptive fields, binocular interaction and functional architecture in cat's visual cortex.** *J Physiol (Lond)* 1962, **160**:106.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

