

Poster presentation

Open Access

Computational model of a modulatory cell type in the feeding network of the snail, *Lymnaea stagnalis*

Dimitris V Vavoulis*^{1,2}, Eugeny S Nikitin², Jianfeng Feng¹, Paul R Benjamin² and György Kemenes²

Address: ¹Ctr. for Scientific Computing, University of Warwick, Coventry, UK and ²Sussex Ctr. for Neuroscience, University of Sussex, Brighton, UK

Email: Dimitris V Vavoulis* - Dimitris.Vavoulis@dcs.warwick.ac.uk

* Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, 8(Suppl 2):P113 doi:10.1186/1471-2202-8-S2-P113

© 2007 Vavoulis et al; licensee BioMed Central Ltd.

Introduction

Realistic mathematical models of single neurons are significant in assessing the contribution of specific ionic conductances to neuronal excitability. This study presents a detailed computational model of the Cerebral Giant Cells (CGCs), a pair of serotonergic neurons in the feeding network of *Lymnaea stagnalis*, which are critical for the expres-

sion of motor behaviour (feeding) and the formation of long-term memory.

Methods

First, we fitted a single-compartment, Hodgkin-Huxley model of the CGCs to two-electrode voltage- and current-clamp data [1] using a combination of linear and non-lin-




Figure 1

Overview of the CGCs model and the contribution of specific currents to neuronal excitability. In **A**, the model has been shifted to the right by 2 msec compared to the biological action potential.

ear least-square fitting techniques. Then, we selectively blocked each ionic current to assess its role in the model, thus mimicking the application of pharmacological agents in the biological neuron.

Results

The model replicates accurately the shape of the action potentials and the tonic firing (~ 0.74 Hz) of the biological neuron (Fig. 1A). A persistent sodium current I_{NaP} and a transient low-threshold calcium current I_{LVA} keep the neuron spontaneously active (Fig. 1Bi, ii). A transient potassium current I_A regulates the interspike interval, while a transient high-threshold calcium current I_{HVA} increases the duration of each spike (Fig. 1Biii, iv). Transient sodium and delayed rectifier potassium currents are responsible for the depolarizing and repolarizing phases of the action potential, as in the classical Hodgkin-Huxley model. The available experimental data [1] are in agreement with these conclusions.

Conclusion

The model we have developed here provides an accurate description of the CGCs at the biophysical level and it is a useful tool for studying the electrical properties of these important modulatory neurons.

Acknowledgements

This research was supported by EPSRC and BBSRC, United Kingdom

References

1. Staras K, Gyori J, Kemenes G: **Voltage-gated ionic currents in an identified modulatory cell type controlling molluscan feeding.** *Eur J Neurosci* 2002, **15**:109-119.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

