

Poster presentation

Open Access

Role of the semi-lunar process in locust jumping

David W Cofer*¹, James Reid², Ying Zhu², Gennady Cymbalyuk³, William J Heitler⁴ and Donald H Edwards¹

Address: ¹Department of Biology, Georgia State University, Atlanta, GA 30303, USA, ²Computer Science, Georgia State University, Atlanta, GA 30303, USA, ³Physics and Astronomy, Georgia State University, Atlanta, GA 30303, USA and ⁴School of Biology, Univ. of St. Andrews, UK

Email: David W Cofer* - dcofer1@student.gsu.edu

* Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, 8(Suppl 2):P12 doi:10.1186/1471-2202-8-S2-P12

© 2007 Cofer et al; licensee BioMed Central Ltd.

The biomechanical and neural components that underlie locust jumping have been extensively studied [1-4]. Previous research suggested that energy for the jump is stored primarily in the extensor apodeme and in the semi-lunar process (SLP) [5], a thickened band of cuticle at the distal end of the tibia. As it has thus far proven impossible to experimentally alter the SLP without rendering a locust unable to jump, it has not been possible to test whether the energy stored in the SLP has a significant impact on the jump, or how that energy is applied during the jump.




Figure 1
Percentage reduction in jump distance without semi-lunar process. Loss of the SLP reduced the distance jumped by approximately 45% across the entire range of extensor tensions tested. Each point is n = 20.

To address problems such as this we have developed a software toolkit, AnimatLab, which allows researchers to build and test virtual organisms. We used this software to build a virtual locust, and then asked how the SLP is utilized during jumping, and how manipulation or removal of the virtual SLP influences jump dynamics (figures 1 and 2). The results show that without the SLP the jump dis-




Figure 2
Power during jump impulse. The power during the early phase of the jump is almost identical, but without the SLP it peaks early and the power from the late phase of the jump is almost entirely missing. This has a significant impact on the jump distance.

tance was reduced by almost half. Further, the simulations were also able to show that loss of the SLP had a significant impact on the final phase of the jump impulse which prevented the full extension of the tibia against the ground. Power for the jump during the initial phase was almost identical between the two cases, but without the SLP the power peaked early and there was a significant difference in the power for the late phase of the jump.

Acknowledgements

This project is supported by GSU Brains & Behavior Program and NIH Grant P20-GM065762.

References

1. Heitler W: **The locust jump.** *J Comp Physiol* 1974, **89**:93-104.
2. Heitler WJ, Burrows M: **The locust jump. II. Neural circuits of the motor programme.** *J Exp Biol* 1977, **66**:221-241.
3. Heitler WJ, Burrows M: **The locust jump. I. The motor programme.** *J Exp Biol* 1977, **66**:203-219.
4. Bennet-Clark HC: **The energetics of the jump of the locust *Schistocerca gregaria*.** *J Exp Biol* 1975, **63**:53-83.
5. Burrows M, Morris G: **The kinematics and neural control of high-speed kicking movements in the locust.** *J Exp Biol* 2001, **204**:3471-3481.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

