



**ORAL PRESENTATION**

**Open Access**

# Cardiac magnetic resonance imaging and gadolinium angiography for neonates and small infants: a 10-year single institutional experience

Sheela Rangamani<sup>1\*</sup>, Joby Varghese<sup>3</sup>, James M Hammel<sup>2</sup>, Scott E Fletcher<sup>1</sup>, Kim F Duncan<sup>2</sup>, David A Danford<sup>1</sup>, Shelby Kutty<sup>1</sup>

From 15th Annual SCMR Scientific Sessions  
Orlando, FL, USA. 2-5 February 2012

## Background

With increasing applications of cardiac magnetic resonance (CMR) for definitive diagnosis of congenital heart disease (CHD) in infants, safety of this technology is of particular interest in this age group. We report our ten-year experience with CMR in neonates and small infants with particular focus on the safety profile and incidence of adverse events.

## Methods

Clinical, anesthesia and nursing records of all patients (pts) <120 days of age who underwent CMR were reviewed. Variables including cardiac diagnosis, study duration, anesthesia type and agents used, prostaglandin E1 (PGE1) dependence and gadolinium (Gd) use were recorded. Serially recorded temperature, systemic saturation, and cardiac rhythm were analyzed. Primary outcome measure was any adverse event (AE) during or <24 hours after the procedure, including minor AE such as hypothermia (axillary temperature  $\leq 95$  F), desaturation (SpO<sub>2</sub> drop  $\geq 10\%$  below baseline) and bradycardia (heart rate  $\leq 100$ /min). Secondary outcome measure was unplanned overnight hospitalization of out-pts.

## Results

Pts (n=143; 74 male, 69 female) had a median age of 7 days (1-117), and 98 were  $\leq 30$  days at the time of CMR. The median weight was 3.4 kg (1.4-6) and body surface area 0.22 m<sup>2</sup> (0.13-0.32). There were 118 (83%) in-pts (108 receiving intensive care) and 25 (17%) out-pts. Indications were assessment of aortic arch (n=57),

complex CHD (n=41), pulmonary veins (n=15), vascular ring (n=8), intracardiac mass (n=8), pulmonary artery (n=7), ventricular volume (n=4), and systemic veins (n=3). CMR was performed using a 1.5-T scanner (Philips), and a commercially available coil. CMR utilized general anesthesia (GA) in 86 pts, deep sedation in 50 and no sedation in 7. Gd angiography was performed in 136 pts. Fifty-nine pts were PGE1 dependent and 39 had a univentricular circulation. Among pts on PGE1, 43 (74%) received GA and 10 (26%) had deep sedation. Twelve pts (8%) had AE, 1 major and 11 minor. Nine of the 12 pts had GA, and 3 had deep sedation. The single major AE was respiratory arrest after chloral hydrate sedation in a neonate (resuscitated without sequelae). Minor AE included desaturations (n=2), hypothermia (n=5), bradycardia (n=2), and bradycardia with hypoxemia (n=2). Incidence of minor AE was 9% for in-pts (vs. 4% for out-pts), and 8% for neonates (vs. 9% in pts  $\geq 30$  days). Incidence of minor AE was similar between PGE1 dependent infants and non-PGE1 group. There were no AE related to Gd angiography. Of 25 out-pts, 5 (20%) were admitted for overnight observation due to desaturations.

## Conclusions

CMR imaging can be accomplished safely in neonates and infants  $\leq 120$  days for a wide range of pre surgical cardiac indications. Incidence of AE was unrelated to pt age, complexity of heart disease, type of anesthesia and PGE1 dependence.

## Funding

None.

<sup>1</sup>University of Nebraska/Creighton University, Joint Division of Pediatric Cardiology, Children's Hospital and Medical Center, Omaha, NE, USA  
Full list of author information is available at the end of the article

#### Author details

<sup>1</sup>University of Nebraska/Creighton University, Joint Division of Pediatric Cardiology, Children's Hospital and Medical Center, Omaha, NE, USA.

<sup>2</sup>Division of Cardiovascular Surgery, University of Nebraska/Creighton University, Children's Hospital and Medical Center, Omaha, NE, USA. <sup>3</sup>Division of Anesthesia, Children's Hospital and Medical Center, Omaha, NE, USA.

Published: 1 February 2012

doi:10.1186/1532-429X-14-S1-O57

**Cite this article as:** Rangamani *et al.*: Cardiac magnetic resonance imaging and gadolinium angiography for neonates and small infants: a 10-year single institutional experience. *Journal of Cardiovascular Magnetic Resonance* 2012 **14**(Suppl 1):O57.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

