

MEETING ABSTRACT

Open Access

Oxidative DNA damage triggers ProNGF-Mediated apoptosis in the striatum of 6-OHDA-treated rats

Tao Wang¹, Pingping Zuo², Piu Chan^{1*}

From 2011 International Conference on Molecular Neurodegeneration
Shanghai, China. 22-24 September 2011

Background

Oxidative DNA damage generated by 6-hydroxydopamine (6-OHDA) may initiate a programmed cell death cascade called apoptosis and plays an important role in the pathogenesis of Parkinson's disease (PD). ProNGF, a precursor of nerve growth factor (NGF), is found to be involved in neuronal apoptosis of neurodegenerative diseases such as PD. Therefore, we have tested the presumption that proNGF-mediated apoptosis might be initiated by oxidative DNA damage in 6-OHDA treated rats.

Method

Enzymatic assay, histochemistry, TUNEL and Western blot were used to investigate profiles of pro-NGF mediated apoptosis after oxidative DNA damage in a unilateral 6-OHDA treated rat model of PD.

Result

A signaling cascade of proNGF-mediated apoptosis definitely occurred in striatum of 6-OHDA lesioned rat under a circumstance of oxidative DNA damage. We confirmed the following changes: 1) activities of antioxidant defense system and levels of malondialdehyde (MDA), 2) MTH1 levels and 8-oxo-2'-deoxyguanosine (8-oxo-dG) accumulation, 3) loss of the TH-positive fiber and dopamine neurons in nigrostriatal system, 4) appearance of apoptotic body and neuronal apoptosis in striatum, 5) expression of proNGF and downstream co-receptor of sortilin and p75NTR, 6) activation status of c-Jun N-terminal kinase (JNK) and its target p53, 7) activation status of the intrinsic apoptotic pathway including cytochrome *c*, caspase 9 and caspase 3, 8) phosphorylation levels of Bad and ratio of Bax/Bcl-2.

Conclusion

These results support that after exposure to the stress stimuli such as oxidative DNA damage induced by 6-OHDA, the destiny of neuronal cells in striatum finally went to apoptosis mediated by proNGF-mediated signaling and it suggests that these profiles might provide some clues for the possible molecular basis of PD etiology.

Author details

¹Department of Neurobiology, Beijing Institute of Geriatrics, Xuanwu Hospital of Capital Medical University, Beijing 100053, China. ²Department of Pharmacology, Institute of Basic Medical Sciences, Peking Union Medical College & Chinese Academy of Medical Sciences, Beijing, China.

Published: 7 February 2012

doi:10.1186/1750-1326-7-S1-S20

Cite this article as: Wang *et al.*: Oxidative DNA damage triggers ProNGF-Mediated apoptosis in the striatum of 6-OHDA-treated rats. *Molecular Neurodegeneration* 2012 **7**(Suppl 1):S20.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Department of Neurobiology, Beijing Institute of Geriatrics, Xuanwu Hospital of Capital Medical University, Beijing 100053, China
Full list of author information is available at the end of the article