

ORAL PRESENTATION

Open Access

Decorrelation of low-frequency neural activity by inhibitory feedback

Tom Tetzlaff^{1*}, Moritz Helias², Gaute T Einevoll¹, Markus Diesmann^{2,3}

From Nineteenth Annual Computational Neuroscience Meeting: CNS*2010 San Antonio, TX, USA. 24-30 July 2010

To correctly judge the functional role of cooperative neural activity it is essential to understand how neural correlations are determined by the structure and dynamics of neural networks. Shared presynaptic input is one of the major sources of correlated synaptic activity in such systems. In the asynchronous state of recurrent neural network models, however, spike correlations are considerably smaller than what one would expect based on the amount of shared presynaptic sources [1,2]. A similar lack of correlations in the spiking activity of neighbouring cortical neurons has been observed experimentally [3]. Recently, it has been pointed out that shared-input correlations can be actively suppressed by the dynamics of recurrent networks [4]. Here, we show that both in networks with purely inhibitory coupling (Fig. 1A) and in those with mixed excitatory-inhibitory coupling (Fig. 1B) this active decorrelation affects

mainly the activity at low frequencies (<20 Hz). High-frequency activity, in contrast, is rather unaffected. Simulations rule out that this phenomenon is the result of refractoriness. By means of a simple linear population-rate model we demonstrate that the effect is essentially explained by inhibitory feedback.

Acknowledgements

Supported by the Research Council of Norway (eVITA, Notur), the Helmholtz Alliance on Systems Biology, the Next-Generation Supercomputer Project of MEXT, EU Grant 15879 (FACETS), DIP F1.2, and BMBF Grant 01GQ0420 to BCCN Freiburg. All network simulations were carried out using NEST (http://www.nest-initiative.org).

Author details

¹Department of Mathematical Sciences and Technology, Norwegian University of Life Sciences, Ås, Norway . ²RIKEN Brain Science Institute, Wako City, Japan . ³Brain and Neural Systems Team, RIKEN Computational Science Research Program, Wako City, Japan.




Figure 1 Population-rate power spectra for an inhibitory (**A**) and a balanced recurrent network (**B**) of leaky integrate-and-fire model neurons (black curves). Grey curves represent population-rate spectra of ensembles of unconnected neurons receiving stationary Poisson input spike trains with the same shared-input structure as in the respective recurrent cases (black curves).

¹Department of Mathematical Sciences and Technology, Norwegian University of Life Sciences, Ås, Norway



^{*} Correspondence: tom.tetzlaff@umb.no

Published: 20 July 2010

References

- Tetzlaff T, Rotter S, Stark E, Abeles M, Aertsen A, Diesmann M: Dependence of neuronal correlations on filter characteristics and marginal spike-train statistics. Neural Comput 2008, 20(9):2133-2184.
- Kriener B, Tetzlaff T, Aertsen A, Diesmann M, Rotter S: Correlations and population dynamics in cortical networks. Neural Comput 2008, 20(9):2185-2226.
- Ecker AS, Berens P, Keliris GA, Bethge M, Logothetis NK, Tolias AS: Decorrelated Neuronal Firing in Cortical Microcircuits. Science 2010, 327(5965):584-587.
- Renart A, de la Rocha J, Bartho P, Hollender L, Parga N, Reyes A, Harris KD: The Asynchronous State in Cortical Circuits. Science 2010, 327(5965):587-590.

doi:10.1186/1471-2202-11-S1-O11

Cite this article as: Tetzlaff *et al.*: Decorrelation of low-frequency neural activity by inhibitory feedback. *BMC Neuroscience* 2010 11(Suppl 1):011.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

