

**The Chondroprotective Actions of a Natural Product are  
Mediated by the Autocrine Activation of IGF-1 Production by  
Human Chondrocytes Despite the Presence of IL-1 $\beta$ .**

Mark J.S. Miller<sup>1</sup>, Salahuddin Ahmed<sup>2</sup>, Paul Bobrowski<sup>3</sup>, Tariq M. Haqqi<sup>2</sup>.

<sup>1</sup>Center for Cardiovascular Sciences, Albany Medical College, Albany, New York

<sup>2</sup> Department of Medicine, Division of Rheumatic Diseases, Case Western  
Reserve University School of Medicine, Cleveland, Ohio

<sup>3</sup> Rainforest Nutritionals, Inc., Raleigh, North Carolina.

Email Addresses

MJSM: [millermj@mail.amc.edu](mailto:millermj@mail.amc.edu)

SA: [salahmed@med.umich.edu](mailto:salahmed@med.umich.edu)

PB: [pbobrowski@rainforest-inc.com](mailto:pbobrowski@rainforest-inc.com)

TMH: [tariq.haqqi@case.edu](mailto:tariq.haqqi@case.edu)

## Abstract

Cartilage loss is a hallmark of arthritis and follows activation of catabolic processes concomitant with a suppression of anabolic pathways like insulin-like growth factor 1 (IGF-1). We hypothesized that two natural products of South American origin, would limit cartilage degradation by respectively suppressing catabolism and activating local IGF-1 anabolic pathways. One extract, derived from cat's claw (*Uncaria guianensis*, vincaria®), is a well-described inhibitor of NF-κB. The other extract, derived from the vegetable *Lepidium meyenii* (RNI 249), possessed an uncertain mechanism of action but with defined ethnomedical applications for fertility and vitality. Human cartilage samples were procured from surgical specimens with consent, and were evaluated either as explants or as primary chondrocytes prepared after enzymatic digestion of cartilage matrix. Assessments included IGF-1 gene expression, IGF-1 production (ELISA), cartilage matrix degradation and nitric oxide (NO) production, under basal conditions and in the presence of IL-1β. RNI 249 enhanced basal IGF-1 mRNA levels in human chondrocytes by 2.7 fold, an effect that was further enhanced to 3.8 fold by co-administration with vincaria. Enhanced basal IGF-1 production by RNI 249 alone and together with vincaria, was confirmed in both explants and in primary chondrocytes (P <0.05). As expected, IL-1β exposure completely silenced IGF-1 production by chondrocytes. However, in the presence of IL-1β both RNI 249 and vincaria protected IGF-1 production in an additive manner (P <0.01) with the combination restoring chondrocyte IGF-1 production to normal levels. Cartilage NO production was dramatically enhanced by IL-1β. Both vincaria and RNI 249 partially attenuated NO production in an additive manner (p< 0.05). IL-1β-induced degradation of cartilage matrix was quantified as glycosaminoglycan release. Individually RNI 249 or vincaria, almost completely prevented this catabolic action of IL-1β. In summary, the identification

of agents that activate the autocrine production of IGF-1 in cartilage, even in the face of suppressive pro-inflammatory, catabolic cytokines like IL-1 $\beta$ , represents a novel therapeutic approach to cartilage biology. When linked to the concomitant functional sequelae, prevention of the catabolic events and the potential for sustained anabolic activity, this natural product combination has significant promise in the treatment of debilitating joint diseases.

**Key Words:**

Arthritis, joint, cartilage, aging, inflammation, IL-1 $\beta$ , IGF-1, anabolic, catabolic, glycosaminoglycan, nitric oxide, complimentary medicine, botanical, natural product, cat's claw, *Uncaria guianensis*, chondrocyte, glucosamine.

## Introduction

Osteoarthritis is a painful and debilitating joint condition that affects hundreds of millions worldwide [1,2]. Despite the prevalence of the disease significant limitations characterize the current therapeutic options. The popular non-steroidal anti-inflammatory class of therapeutics (NSAIDs), which block cyclo-oxygenase (COX), provide symptomatic relief but do not abrogate the underlying disease process [3]. Indeed, it is well appreciated that cartilage destruction can proceed unabated despite suppression of inflammation [4]. To this dilemma, the COX-2 specific class of NSAIDs has revealed an increase in the risk for cardiovascular and heart disease [5,6], although their design was an attempt to reduce complications associated with non-specific COX inhibitors [7]. Thus the quest to develop new therapeutic entities has taken on greater impetus and yet additional uncertainty.

Patients often supplement pharmaceutical strategies for managing arthritis with complimentary medicines. These may include acupuncture [8], nutraceuticals [9-11], and botanicals [12,13]. Glucosamine and chondroitin based products dominates the nutraceutical approach to arthritis. Glucosamine and chondroitin are structural elements of cartilage and matrix and their proposed therapeutic benefits center on the assumption that ingestion of these matrix elements, despite their poor absorption [14,15], will replace matrix that is lost as a result of the catabolic process. Not surprisingly large amounts are required and the onset of action with this approach is on the order of months [16-19].

Botanicals, especially those with redox-based actions, hold promise in the treatment of chronic inflammation [20-25]. Green tea catechins, especially epigallocatechin gallate (EGCG), have been shown to limit human cartilage degradation in vitro [20,26] and maintain joint architecture in the CIA animal model [25]. This anti-inflammatory action is

thought to be the result of inhibition of transcriptional events, particularly prevention of NF- $\kappa$ B activation by cytokines and oxidants. NF- $\kappa$ B is a critical transcription factor in chronic inflammation and is a desirable target for new therapeutics, including pharmaceutical development, as it regulates numerous genes that contribute to the inflammatory process [27-29]. Of particular note for joint dysfunction, NF- $\kappa$ B regulates the production of matrix metalloproteases (MMPs) by chondrocytes [30]. During inflammation or injury chondrocytes release MMPs, which in turn degrade the cartilage matrix, releasing glycosaminoglycans.

While the developmental pipeline for new therapeutics has focused on NSAIDs and approaches to alleviate joint destruction, there are few approaches that also address restoration or activation of anabolic, repair pathways. It has been well appreciated that inflammation and infection silences the expression of repair pathways like IGF-1 [31,32]. Concomitant with a suppression of local, autocrine production of IGF-1 during inflammation, is an uncoupling of the signal transduction pathways for exogenous Growth Hormone (GH) and IGF-1 [33-35]. This combination of lost tissue responsiveness to circulating levels of anabolic factors and compromised local production results in a sustained loss of anabolic tone. For joints, this may include supporting musculature as well as cartilage and bone [36]. This problem is more apparent for the elderly where these events are superimposed on an age-related loss in IGF-1 activity. For these reasons, Cappola et al [37] observed that those elderly individuals with a combination of high IL-6 levels and low IGF-1 levels are at the greatest risk for enhanced morbidity and mortality.

These factors also relate to the increasing incidence of osteoarthritis and joint dysfunction with age [38] and the reduced growth associated with pediatric chronic inflammatory bowel disease, where anti-IL-6 antibodies restored linear growth and IGF-1 levels independent of nutrition [39]. Indeed, there appears to be excellent evidence that growth restriction with chronic inflammation, including arthritis, is the result of suppression of IGF-1 production and activity, independent of GH, and in response to IL-6 [40,41]. Supplemental IGF-1 restored linear growth in these models, linking the suppression of IGF-1 to compromised anabolic activity [42].

In the present study we hypothesized that a combination of a cat's claw (*Uncaria guianensis*) extract devoid of immunostimulatory oxindole alkaloids [43] and an extract of the Andean vegetable *Lepidium meyenii*, may provide the desired therapeutic innovation for compromised anabolic tone. The rationale for this hypothesis centered on our previous work with *Uncaria guianensis* extract vincaria, as a potent inhibitor of NF- $\kappa$ B, TNF $\alpha$  and associated anti-inflammatory and cytoprotective actions [24, 43-45]. Additionally, vincaria has successfully completed a small trial for osteoarthritis of the knee demonstrating a rapid reduction in pain and improved function [46].

*Lepidium meyenii* extracts had previously been demonstrated to have pro-fertility actions [47-50], an application that is not intuitive for treating arthritis. The pro-fertility actions of *Lepidium meyenii* could not be explained on the basis of an augmentation of sex steroid or gonadotrophin pathways [47,48] and the mechanism underlying these benefits has remained elusive until now. However, based on this ethnomedical knowledge and our own observations that these extracts were anabolic in farmed fish (unpublished reports) whose growth is critically dependent on IGF-1, we hypothesized that the central

mechanism of action for *Lepidium meyenii* extracts was the autocrine activation of IGF-1. If our hypothesis was to be validated then this would rapidly open a new approach to the management of arthritis.

The present study was designed to address these issues with a focus on the functional consequences of maintaining IGF-1 gene activity and production in human cartilage explants. The study had goals of determining if RNI 249 could activate human chondrocyte IGF-1 gene expression and maintain IGF-1 production in the face of pro-inflammatory signals, IL-1 $\beta$ , that normally silence the IGF-1 gene.

## **Materials and Methods**

### ***Reagents***

Tissue culture medium and related reagents were purchased from either Mediatech (Herndon, VA) or InVitrogen (Carlsbad, CA). Recombinant human IL-1 $\beta$  was purchased from R&D Systems (St Paul, MN), and other chemicals were purchased from Sigma-Aldrich (Saint Louis, MO). The extracts, vincaria (RN180) or RNI 249, and their combination (Reparagen™) were supplied by Rainforest Nutritionals, Inc and were dissolved in water and filtered through a 0.45  $\mu$ m filter under vacuum prior to use. Vincaria is an alkaloid depleted water based extract of *Uncaria guianensis* that is standardized for antioxidant activity (DPPH radical quenching) and an oxindole alkaloid content of less than 0.1 mg/g as determined by HPLC. RNI 249 consists of a polar extract of *Lepidium meyenii* that is standardized for its DPPH quenching activity prior to combining with an inert stabilizing material.

### ***Human Chondrocytes Culture and Cartilage Explants***

Human OA cartilage samples were procured through the Cooperative Human Tissue Network and with prior approval of the Institutional Review Board of University Hospitals of Cleveland. Full-thickness cartilage slices (20-25 mg) were dissected from the cartilage using sterile scalpel blade (Feather Safety Razor Co., Japan). Four to five cartilage pieces (approximately equal in size and weight) were transferred to each well of a 24-well, flat bottom plate (Nunc, Denmark) containing DMEM:F-12 (1:1) supplemented with antibiotics and 10% FCS. Chondrocytes were prepared by the enzymatic digestion of knee cartilage as previously described [20,23,26] and maintained in DMEM:F12 (Mediatech, Herndon, VA) supplemented with 10% FBS. The cartilage explants were treated with IL-1 $\beta$  alone or with IL-1 $\beta$  + Vincaria or RNI 249, alone or in combination for 72 hrs. Explants cultured in the absence of IL-1 $\beta$ , vincaria or RNI 249, were used as controls. Additionally, the actions of RNI 249 and vincaria singularly and together were examined independently of IL-1 $\beta$  exposure (5 ng/ml). Total glycosaminoglycan present in the culture supernatant was estimated as described below.

### ***Quantitative RT-PCR.***

Total cytoplasmic RNA was prepared from human chondrocytes using a commercially available kit according to the instructions of the manufacturer (Qiagen, Valencia, CA). Real time quantitative RT-PCR with internal fluorescent hybridization probes was performed as previously described [21] and the IGF-1 gene expression was quantified using a commercially available Gene Expression Assay kit (Applied Biosystems, CA). Expression of IGF-1 was normalized to  $\beta$ -actin mRNA expression, and the results were expressed as fold induction relative to controls.

### ***IGF-1 ELISA***

Human IGF-1 level in culture or explant media was quantified using a commercially available Human IGF-1 ELISA kit (R & D Systems) per manufacturer's directions.

### ***Quantitation of Glycosaminoglycans.***

At the end of culture period, the culture medium was collected from each group. A 50  $\mu$ l aliquot of the collected supernatant from each sample was utilized to estimate the total glycosaminoglycan concentration by a colorimetric method employing DMMB as previously described [52]. Color intensity was read spectrophotometrically at 535 nm using the Lambda 25 spectrophotometer (Perkin-Elmer, CT) and the values were derived from a standard curve prepared using different concentrations of chondroitin sulfate. Results are expressed as micrograms of glycosaminoglycan released per milligram of cartilage tissue.

### ***Nitric Oxide Production***

Levels of nitrate/nitrite were measured in cartilage explant and chondrocyte culture media by using the Griess reaction (colorimetric assay) after conversion of the nitrate to nitrite using a commercially available kit (Molecular Probes, OR).

### ***Statistical Analysis***

Each experiment was repeated to ensure reproducibility of the data using cartilage samples from matched donors. Data was analyzed using the software package, InStat®, and the analysis included one way ANOVA followed by an appropriate post hoc test (Dunnett's). Values are expressed as mean  $\pm$  SEM. Differences were considered significant at  $P < 0.05$ .

## Results

### ***IGF-1 Expression and Production by Human Cartilage***

Direct activation of IGF-1 gene expression was evaluated in human chondrocytes by real time RT-PCR. RNI 249 (50µg/ml) as well as vincaria (10 µg/ml) when added to the culture media for 48hr resulted in a significant increase in the expression of IGF-1 (Figure 1), relative to the control, β-actin (P<0.05). Furthermore, when RNI 249 and vincaria were combined, these effects were additive resulting in a nearly 4-fold increase in IGF-1 gene expression, which was significantly different from RNI 249 or vincaria alone, as well as basal expression (P<0.001).

The potential for RNI 249 to stimulate chondrocyte IGF-1 production was assessed by measurement of IGF-1 in the culture media of cartilage explants. Explants were exposed to RNI 249 at 10 and 50 µg/ml for 48 hours at which time the media was removed for ELISA determination of human IGF-1 levels (Figure 2). RNI 249 treatment resulted in a dose-dependent increase in IGF-1 levels that were significant at the 50 µg/ml dose (P<0.05). RNI 249 represents the combination of a proprietary *Lepidium meyenii* extract with inert stabilizers as it was noted that the extract alone had a reduced shelf-life, which is corrected in the RNI 249 form.

IGF-1 one production from human cartilage explants was also enhanced 67% by vincaria alone (10 µg/ml, P<0.05), consistent with the stimulatory effects of vincaria treatment on IGF-1 gene expression indicated in Figure 1.

### ***Effects of IL-1 $\beta$ on IGF-1 Production***

While RNI 249 stimulates basal IGF-1 gene expression and production in human cartilage, it was necessary to determine if this action was retained in the presence of IL-1 $\beta$ . Pro-inflammatory cytokines like TNF $\alpha$  and IL-1 $\beta$  are known to silence the expression of the IGF-1 gene and shutdown production [31,32,41]. This ability of IL-1 $\beta$  was confirmed in cartilage explants where IL-1 $\beta$  (5 ng/ml) exposed explants displayed a complete suppression of IGF-1 production (Figure 3). Treatment with either vincaria (10  $\mu$ g/ml) or RNI 249 (50  $\mu$ g/ml) partially restored IGF-1 production ( $P < 0.01$ ) and the combination fully restored IGF-1 levels to baseline despite the presence of IL-1 $\beta$ .

### ***IL-1 $\beta$ and GAG Release***

Exposure of human cartilage explants to IL-1 $\beta$  resulted in an increase release of glycosaminoglycans (GAG) into the media (Figure 4,  $P < 0.05$ ). When RNI 249 was added to IL-1 $\beta$  treated explants there was a dose-dependent attenuation of GAG release. Explants that received the 50  $\mu$ g/ml dose of RNI 249 were not statistically significant from controls and were significantly improved over the 10  $\mu$ g/ml dose and IL-1 $\beta$  alone ( $P < 0.01$ ).

Vincaria (10 $\mu$ g/ml) also demonstrated an ability to suppress IL-1 $\beta$  induced GAG release (control =  $0.141 \pm 0.009$   $\mu$ g/mg; IL-1 alone =  $0.223 \pm 0.010$   $\mu$ g/mg; IL-1 $\beta$  + vincaria =  $0.175 \pm 0.010$   $\mu$ g/mg;  $P < 0.01$  for IL-1 vs. control or IL-1 + vincaria,  $n=3$ ). Potential additive effects associated with the combination of RNI 249 and vincaria was not tested as the individual components provided essentially full protection as they were not significantly different from control untreated explants.

### ***IL-1 $\beta$ and Chondrocyte NO Production***

Media nitrite levels in cartilage explants and chondrocyte cultures were used as an index of nitric oxide production. Both RNI 249 (50  $\mu\text{g/ml}$ ) and vincaria (10 $\mu\text{g/ml}$ ) lowered basal nitrite levels (Figure 5) suggesting that there was a mild inherent activation of inducible nitric oxide synthase in these surgical samples. Nevertheless, IL-1 $\beta$  exposure resulted in a marked increase in nitrite levels consistent with an activation of inducible nitric oxide synthase. This response was partially attenuated by vincaria and RNI 249 alone, with additive effects upon co-treatment ( $P < 0.01$ ).

### **Discussion**

Circulating levels of IGF-1 primarily reflect the hepatic production under the influence of growth hormone (GH). While reduced production of GH and IGF-1 are associated with aging and joint dysfunction, poor mobility and increased mortality [36,37,53], the system can be further uncoupled at the level of signal transduction. Both the ability of GH to stimulate IGF-1 formation, and the ability of IGF-1 to activate anabolism can be blocked by signal transduction that fails to lead to transcriptional events [34-36]. Increased production of IGF binding proteins that scavenge IGF also contribute to the overall functional tone of the system [31,42,54-56].

While there are a number of factors including cytokines and poor nutrition, that are thought to contribute to a disruption in IGF-1 functionality [57], the present observations are the first to report that autocrine activation of IGF-1 gene expression in target tissues is a possible solution to these problems. We confirmed in human cartilage explants what has been observed in other tissues that pro-inflammatory cytokines silence IGF-1 production [31,32,35]. In the present case this was achieved with IL-1 $\beta$ , a major

determinant of chondrocyte activation and cartilage destruction [3,4,33,58]. However, this is the first report that IGF-1 production can be maintained in the face of these otherwise completely suppressive signals.

The ability of RNI 249, alone and in combination with vincaria, to maintain IGF-1 levels despite the presence of IL-1 $\beta$  has the potential to not only limit cartilage destruction, as was confirmed here with blockade of GAG release, but also to evoke anabolic actions and repair the joint. The present study was not focused on a clear demonstration of an anabolic action but this is a reasonable outcome based on the known actions of IGF-1 [58-63].

It is not known how these botanical extracts achieve these results. We do propose, based on previous observations that vincaria is effective by limiting the inhibitory actions of IL-1 $\beta$  or the suppressive tone of other pro-inflammatory agents. Vincaria is a remarkably potent redox-based inhibitor of NF- $\kappa$ B and has an IC<sub>50</sub> for inhibiting LPS-induced TNF $\alpha$  production of 10ng/ml [24, 43, 45, 46]. Thus we propose that vincaria operates by blocking the inhibitory actions of inflammatory mediators on basal IGF-1 production rather than a direct activation in a manner that is similar to that described by Haqqi and colleagues [20,21].

On the other hand, RNI 249, which is a far weaker antioxidant than vincaria [39,40,58], and its parent vegetable source, *Lepidium meyenii*, and as such is unlikely to operate via redox-based actions. *Lepidium meyenii* has primarily been researched for its ability to promote fertility in males and females. The breadth of *Lepidium meyenii*'s actions on fertility is impressive and range from increased fetal growth, reduced miscarriage rates,

improvements in conception rates and increased semen volumes and sperm counts [47-51]. It is well documented that *Lepidium meyenii* does not affect the levels of gonadotrophins, sex steroids or prolactin [47-48], and hence the mechanism of action for these pro-fertility effects has remained elusive to date. As IGF-1 is a critical determinant of fertility and fetal development, we propose that the actions of *Lepidium meyenii* are dictated by a central action of the direct, local activation of IGF-1 production in target tissues independently of growth hormone. What is of critical interest is that this action can be observed in the presence of IL-1 $\beta$ , a potent inflammatory agent. Furthermore, when RNI 249 is combined with the anti-inflammatory botanical vincaria, there are additive effects, fully restoring balance where previously the forces were tipped towards catabolism and inflammation.

RNI 249 alone, as well as vincaria, was able to block the increased release of GAG from cartilage explants by IL-1 $\beta$ . This action of vincaria had been noted previously when combined with a mineral supplement [23], and similar effects have also been observed with potent redox-based anti-inflammatories – green tea epigallocatechin gallate [21] and anthocyanidin enriched pomegranate fruit extract [22]. These redox based natural agents appear to limit cartilage breakdown by blocking the expression, production and release of matrix metalloproteases by activated chondrocytes via a NF- $\kappa$ B and MAPKinase dependent pathway.

In the case of RNI 249 this limitation of MMP-mediated events is unlikely to be due to a direct suppression of transcription factors like NF- $\kappa$ B, as has been demonstrated by the other botanicals, but rather a consequence of the autocrine activation of IGF-1. IGF-1 and Th2 cytokines can limit the production and actions of pro-inflammatory signals [61-

63]. Administration of exogenous IGF-1 has been shown to confer protection to joints by silencing pro-inflammatory genes that promote cartilage destruction and joint remodeling [54,61-63]. It was of interest to note that nitric oxide production was less susceptible to normalization than was GAG release. This follows a pattern seen with other botanical extracts or nutraceuticals, which note reduced but not complete blockade of IL-1 $\beta$  induced nitric oxide production [21-23]. This suggests that these different pathways vary in their responsiveness to these interventions, and this may impact the clinical response in terms of individual variability and disease targeting. While incomplete, the attenuation of nitric oxide production reflects the transcriptional regulatory actions of this technology as well as enhancing the effectiveness of the IGF-1 axis. Studer [65] recently described that nitric oxide modifies IGF-1 receptor kinase probably via the formation of nitrosothiols. This structural modification resulted in altered functionality with NO blockade enhancing IGF-1 mediated proteoglycan synthesis. Thus the present approach which leverages systems that suppress NO formation as well as directly raising IGF-1 production will likely yield enhanced restorative, anabolic actions within chondrocytes. How this would be played out in the setting of autocrine activation of IGF-1 production versus exogenous IGF-1 administration or circulating IGF-1 is less clear.

Similarly, either exogenous IGF-1 or a c-Jun N-Terminal kinase (JNK) inhibitor has been shown to protect proteoglycan synthesis following brief thermal stress in cartilage explants [61]. This was in part mediated by a protective action against untoward apoptosis and draws similarities to the anti-apoptotic actions and transcriptional inhibitory actions of vincaria [224,43,44]. This offers further rationalization for such a combination as a therapeutic entity.

There have numerous and diverse approaches employed to enhance the anabolic actions of IGF-1 ranging from increasing production or independently enhancing functionality. Haupt et al [55] using viral gene transfer studies to synoviocytes corrected articular cartilage degradation in vitro. They observed that by combining IGF-1 with an IL-1 receptor antagonist protein they could both increase proteoglycan production and decrease catabolism via reduced MMP synthesis. They proposed that a combination approach of combining anabolic growth factors with catabolic blockers to prevent matrix degradation would be ideal for healing destructive joint diseases. This is conceptually similar to the approach used in the present study – enhanced autocrine production of IGF-1 and suppression of inhibitory cytokine tone. The contrasting mechanistic approaches to the cartilage biology by vincaria and RNI 249 may contribute to the observations that vincaria and RNI 249 activities were additive (suppression of NO release, stimulation of IGF-1 production), although it is acknowledged that full dose-response curves were not performed and so this interpretation cannot be affirmed with certainty.

Madry et al [63] demonstrated that IGF-1 gene transfer was effective healing for traumatic cartilage repair. Using an in vivo model of cartilage loss in rabbits, Tuncel et al [62] noted that collagen sponges impregnated with recombinant IGF-1 accelerated cartilage restoration. Another approach was described by De Ceuninck et al [54], who noted that disruption of IGF-1 binding to the inhibitory binding proteins via small molecules, resulting in the release of free IGF-1, restores the anabolic response of human chondrocytes. This suggests that a critical regulatory site of IGF-1 bioactivity is the inactivation of circulating IGF-1 by binding proteins. With the appreciation that increased levels of IGF-1 binding proteins have been observed in osteoarthritic cartilage [56] one needs to consider if the optimal approach is to raise circulating IGF-1 levels or

alternatively promote local production in an autocrine manner where the potential for binding protein inactivation is reduced.

While these observations collectively demonstrate proof of principle that IGF-1 can promote healing and joint restoration, and that a multitude of potential approaches to the problem are possible, few are likely to have an immediate impact on health care. By contrast, the combination of the natural products RNI 249 and vincaria has already been commercialized in the USA and Canada under the name “Reparagen” and so it can be considered an immediate and current option in contrast to the elegant but commercially distant approaches of gene transfer or other pharmacological approaches.

A clear limitation of the present study is that it has been performed in vitro, albeit with human cartilage specimens, and therefore it would be appropriate to hold reservations that the present observations may not be translated into the in vivo setting or effective with oral intake. However, we consider that outcomes as being unlikely. In the case of vincaria it has already been shown that it is effective in treating osteoarthritis of the knee in humans with oral administration at the remarkably low dose of only 100mg/day [46]. Additionally, vincaria has displayed oral bioactivity in animal models of inflammation [24,43]. RNI 249 is also likely to be orally active in terms of enhanced IGF-1 production as it is orally active for anabolic actions in farmed fish (unpublished observations) and *Lepidium meyenii* and related extracts have well defined effects in pregnancy and male fertility upon oral administration [47-50]. However, an osteoarthritis orientated clinical trial would address these issues and place the present innovation in an appropriate perspective where one could also address potential benefits to the joint as a whole, including skeletal muscle and bone.

## **Conclusion**

Joint inflammation and destruction are not well correlated. Often therapies provide symptomatic relief but fail to restore architecture. The present observations with a novel South American medicinal plant combination that simultaneously addresses catabolic and anabolic activities, suggests that it is possible to directly activate local, chondrocyte IGF-1 production, even in the presence of IL-1 $\beta$ . This action has the potential to provide a therapeutic approach that not only limits joint inflammation and destruction but to promote joint restoration.

## **List of Abbreviations**

MMP matrix metalloproteinase

IGF-1 insulin-like growth factor-1

JNK c-Jun N-Terminal kinase

NF- $\kappa$ B nuclear factor kappa B

IL-1 $\beta$  interleukin 1 beta

IL-6 interleukin 6

TNF $\alpha$  tumor necrosis factor alpha

LPS lipopolysaccharide

RT-PCR reverse transcriptase polymerase chain reaction

GAG glycosaminoglycan

NSAID non-steroidal anti-inflammatory drug

COX cyclo-oxygenase

GH growth hormone

IC<sub>50</sub> inhibitory concentration 50 percent

### **Competing Interests**

MJSM has a financial interest in Rainforest Nutritionals, Inc

PB has a financial interest in Rainforest Nutritionals, Inc

A SBIR grant award to Rainforest Nutritionals, Inc was used to sponsor the research performed by SA and TMH in this study.

### **Authors Contributions**

MJSM contributed to the design of the study and was the lead author of the manuscript.

SA was the key individual that conducted the studies.

PB contributed to study design and manuscript preparation.

TMH oversaw the performance of the study and contributed to its design and manuscript preparation.

### **Acknowledgements**

This work was sponsored by a Small Business Innovation Research Award NIH R43 AG024733-01, from the National Institute on Aging, National Institutes of Health, USA

### **References**

1. Murray CJL, Lopez AD: *The Global Burden of Disease*. Boston: Harvard University Press, 1996.

2. Yellin E, Callahan LF. **The economic cost and social and psychological impact of musculoskeletal conditions.** *Arthritis Rheumatism* 1995, **38**: 1351-1362.
3. Kraan PM, van den Berg WM. **Anabolic and destructive mediators in osteoarthritis.** *Curr Opin Nutr Metab Care* 2000, **3**: 205-211.
4. van den Berg WB. **Joint inflammation and cartilage destruction may occur uncoupled.** *Springer Semin Immunopathol* 1998, **20**: 149-164.
5. Graham DJ, Campen D, Hui R, Spence M, Cheetham C, Levy G, Shoor S, Ray WA. **Risk of acute myocardial infarction and sudden cardiac death in patients treated with cyclo-oxygenase 2 selective and non-selective non-steroidal anti-inflammatory drugs: nested case-control study.** *Lancet.* 2005, **365**: 475-81.
6. Kimmel SE, Berlin JA, Reilly M, Jaskowiak J, Kishel L, Chittams J, Strom BL. **Patients exposed to rofecoxib and celecoxib have different odds of nonfatal myocardial infarction.** *Ann Intern Med.* 2005, **142**: 157-64.
7. Dickman A, Ellershaw J. **NSAIDs: gastroprotection or selective COX-2 inhibitor?** *Palliat Med* 2004, **18**: 275-86.
8. Berman BM, Lao L, Langenberg P, Lee WL, Gilpin AM, Hochberg MC. **Effectiveness of acupuncture as adjunctive therapy in osteoarthritis of the knee: a randomized, controlled trial.** *Ann Intern Med* 2004, **141**: 901-10.
9. Noack W, Fischer M, Forster KK, Rovarti LC, Setnikar A. **Glucosamine sulfate in osteoarthritis of the knee.** *Osteoarthritis Cartilage* 1994, **2**: 61-69.
10. Walter-Bone K. **'Natural remedies' in the treatment of osteoarthritis.** *Drugs Aging* 2003, **20**: 517-526.

11. Hesslink R, Armstrong D, Nagendra MV, Sreevatsan S, Barathur R. **Cetylated fatty acids improve knee function in patients with osteoarthritis.** *J Rheumatol* 2002, **29**: 1708-1712.
12. Tao X, Lipsky PE. **The Chinese anti-inflammatory and immunosuppressive herbal remedy Tripterygium wilfordii Hook F.** *Rheum Dis Clin North Am* 2000, **57**: 1221-1227.
13. Aggarwal BB, Shishodia S. **Suppression of the nuclear factor- $\kappa$ B activation pathway by spice-derived phytochemicals: reasoning for seasoning.** *Ann NY Acad Sci* 2004, **1030**: 434-441.
14. Biggee BA, Blinn CM, McAlindon TE, Nuite M, Silbert JE. **Low levels of human serum glucosamine after ingestion of glucosamine sulphate relative to capability for peripheral effectiveness.** *Ann Rheumatol* 2005: online pub. Aug 3.
15. McAlindon TE and Biggee BA. **Nutritional factors and osteoarthritis: recent developments.** *Curr Opin Rheumatol* 2005, **17**: 647-652.
16. Cibere J, Kopec JA, Thorne A, Singer J, Canvin J, Robinson DB, Pope J, Hong P, Grant E, Esdaile JM. **Randomized, double-blind, placebo-controlled glucosamine discontinuation trial in knee osteoarthritis.** *Arthritis Rheumat* 2004, **51**: 738-745.
17. Reginster JY, Deroisy R, Rovati LC, Lee RL, Lejeune E, Bruyere O, Giacovelli G, Henrotin Y, Dacre JE, Gossett C. **Long-term effects of glucosamine sulphate on osteoarthritis progression: a randomised, placebo-controlled clinical trial.** *Lancet* 2001, **357**: 251-256.
18. Pavelka K, Gatterova J, Olejarova M, Machacek S, Giacovelli C, Rovati LC. **Glucosamine sulphate use and delay of progression of knee osteoarthritis:**

- a 3-year, randomized, placebo-controlled, double-blind study.** *Arch Intern Med* 2002, **162**: 2113-2123.
19. McAlindon TE, Formica M, LaValley M, Lehmer M, Kabbara K. **Effectiveness of glucosamine for symptoms of knee osteoarthritis: results from an internet-based randomized double-blind controlled trial.** *Am J Med* 2004, **117**: 643-649.
20. Ahmed S, Rahman A, Hasnain A, Lalonde M, Goldberg VM, Haqqi TM. **Green tea polyphenol epigallocatechin-3-gallate inhibits the IL-1 $\beta$ -induced activity and expression of cyclo-oxygenase-2 and nitric oxide synthase-2 activity in human chondrocytes.** *Free Radical Biology Med* 2002, **33**: 1097-1105.
21. Ahmed S, Anuntiyo J, Malesud CJ, Haqqi TM. **Biological basis for the use of botanicals in osteoarthritis & rheumatoid arthritis: a review.** *ECAM* 2005, **2**: 301-308.
22. Ahmed S, Wang N, Hafeez BB, Cheruvu VK, Haqqi TM. **Punica granatum L. extract inhibits IL-1 $\beta$ -Induced expression of matrix metalloproteinases by inhibiting the activation of MAP kinases and NF- $\kappa$ B in human chondrocytes in vitro.** *J Nutr* 2005, **135**: 2096 – 2102.
23. Miller MJS, Ahmed S, Bobrowski P, Haqqi TM. **Suppression of human cartilage degradation and chondrocyte activation by a unique mineral supplement (SierraSil™) and a cat's claw extract, vincaria®.** *J Am Nutraceut Assoc* 2004, **7**: 32-39.
24. Sandoval-Chacon M, Thompson JH, Liu X, Mannick EE, Sadowska-Krowicka H, Charbonnet R, Clark DA, Miller MJS. **Anti-inflammatory actions of cat's claw: the role of NF- $\kappa$ B.** *Alimentary Pharmacol Ther* 1998, **12**: 1279-1289.

25. Haqqi TM, Anthony DD, Gupta S, Ahmed N, Lee M-S, Kumar GK, Mukhtar H. **Prevention of collagen induced arthritis in mice by a phenolic fraction of green tea.** *Proc Natl Acad Sci USA* 1999, **96**: 4525-4529.
26. Singh R, Ahmed S, Malemud CJ, Goldberg VM, Haqqi TM. **Epigallocatechin-3-gallate selectively inhibits interleukin-1 $\beta$ -induced activation of mitogen activated protein kinase subgroup c-Jun N-terminal kinase in human osteoarthritis chondrocytes.** *J Orthopaedic Res* 2003, **21**: 102-109.
27. Marok R, Winyard PG, Coumbe A, Kus ML, Gaffney K, Blades S, Mapp PI, Morris CJ, Blake DR, Kaltschmidt C, Baeuerle PA. **Activation of the nuclear transcription factor- $\kappa$ B in human inflamed synovium.** *Arthritis Rheum* 1996, **39**: 583-591.
28. Winyard PG, Blake DR. **Antioxidants, redox-regulated transcription factors.** *Adv Pharmacol* 1997, **18**: 403-421.
29. Lohmander LS. **What can we do about osteoarthritis?** *Arthritis Res* 2000, **2**: 95-100.
30. Mix KS, Mengshol JA, Benbow U, Vicnenti MP, Sporn MB, Brinkerhoff CE. **A synthetic triterpenoid selectively inhibits the induction of matrix metalloproteinases 1 and 13 by inflammatory cytokines.** *Arthritis Rheum* 2001, **44**: 1096-1104.
31. Fernandez-Celemin L, Pasko N, Blomart V, Thiessen J-P. **Inhibition of muscle insulin-like growth factor 1 expression by tumor necrosis factor-alpha.** *Am J Physiol Endocrinol Metab* 2002, **283**: E1279-E1290.
32. Soto L, Martin AI, Vara E, Lopez-Calderon A. **Cyclosporin A treatment is able to revert the decrease in circulating GH and IGF-1 and increase in IGFBPs induced by adjuvant arthritis.** *Horm Metab Res* 2001, **33**: 590-595.

33. Loeser RF, Carlson CS, Del Carlo M, Cole A. **Detection of nitrotyrosine in aging and osteoarthritic cartilage. Correlation of oxidative damage with presence of IL-1B and with chondrocyte resistance to insulin like-growth factor 1.** *Arthritis Rheum* 2002, **46**: 2349-2357.
34. Holzenberger M. **The role of insulin-like signalling in the regulation of ageing.** *Horm Res* 2004; **62 Suppl 1**: 89-92.
35. Strle K, Broussard SR, McCusker RH, Shen WH, Johnson RW, Freund GG, Dantzer R, Kelley KW. **Proinflammatory cytokine impairment of insulin-like growth factor 1-induced protein synthesis in skeletal muscle myoblasts requires ceramide.** *Endocrinology* 2004, **145**: 4592-4602.
36. Barbieri M, Ferrucci L, Ragno E, Corsi A, Bandinelli S, Bonafe M, Oliveri F, Giovagneti S, Franscheschi C, Guralnik JM, Paolisso G. **Chronic inflammation and the effects of IGF-1 on muscle strength and power in older persons.** *Am J Physiol Endocrinol Metab* 2003, **284**: E481-E487.
37. Cappola AR, Xue QL, Ferrucci L, Guralnik JM, Volpato S, Fried LP. **Insulin-like growth factor and interleukin-6 contribute synergistically to disability and mortality in older women.** *J Clin Endocrinol Metab* 2003, **88**: 2019-2025.
38. Ponzer S, Tidermark J, Brismar K, Soderqvist A, Cederholm T. **Nutritional status, insulin-like growth factor-1 and quality of life in elderly women with hip fractures.** *Clin Nutr* 1999, **18**: 241-246.
39. Sawczenko A, Azooz O, Paraszczuk J, Idestrom M, Savage MO, Ballinger AB, Sanderson IR. **Intestinal inflammation-induced growth restriction acts through IL-6 in rats and depends on the – 174 IL-6 G/C polymorphism in children.** *Proc Natl Acad Sci (USA)* 2005, **102**: 13260-13265.
40. De Benedetti F, Meazza C, Oliveri M, Pignatti P, Vivarelli M, Alonzi T, Fattori E, Garrone S, Barreca A, Martini A. **Effect of IL-6 on IGF binding protein-3: a**

- study in IL-6 transgenic mice and in patients with systemic juvenile idiopathic arthritis.** *Endocrinology* 2001, **142**: 4818-26.
41. De Benedetti F, Alonzi T, Moretta A, Lazzaro D, Costa P, Poli V, Martini A, Ciliberto G, Fattori E. **Interleukin 6 causes growth impairment in transgenic mice through a decrease in insulin-like growth factor-I. A model for stunted growth in children with chronic inflammation.** *J Clin Invest* 1997, **99**: 643-50.
42. Ballinger AB, Azooz O, El-Haj T, Poole S, Farthing MJ. **Growth failure occurs through a decrease in insulin-like growth factor 1 which is independent of undernutrition in a rat model of colitis.** *Gut* 2000, **46**: 694-700.
43. Sandoval M, Okuhama NN, Zhang X-J, Condezo LA, Lao J, Angeles FM, Bobrowski P, Miller MJS. **Anti-inflammatory and antioxidant activities of cat's claw (*Uncaria tomentosa* and *Uncaria guianensis*) are independent of their alkaloid content.** *Phytomedicine* 2002, **9**: 325-337.
44. Miller MJS, Angeles FM, Reuter BK, Bobrowski P, Sandoval M. **Dietary antioxidants protect gut epithelial cells from oxidant induced apoptosis.** *BMC Compl Altern Med* 2001, **1**: 11.
45. Sandoval M, Charbonnet RM, Okuhama N, Roberts J, Krenova Z, Trentacosti AM, Miller MJS. **Cat's claw inhibits TNF $\alpha$  production and scavenges free radicals: role in cytoprotection.** *Free Radical Biol Med* 2000, **29**: 71-78.
46. Piscocoya J, Rodriuez Z, Bustamante SA, Okuhama NN, Miller MJS, Sandoval M. **Efficacy and safety of freeze-dried cat's claw in osteoarthritis of the knee: mechanisms of action of the species *Uncaria guianensis*.** *Inflamm Res* 2001, **50**: 442-448.
47. Gonzales GF, Cordova A, Vega K, Chiung A, Villena A. **Effect of *Lepidium meyenii* (Maca), a root with aphrodisiac and fertility-enhancing properties,**

- on serum reproductive hormone levels in adult healthy men. *J Endocrinol* 2003, **176**: 163-168.
48. Gonzales GF, Cordova A, Vega K, Chung A, Villena A, Gonez C, Castillo S. **Effect of *Lepidium meyenii* (MACA) on sexual desire and its absent relationship with serum testosterone levels in adult healthy men.** *Andrologia* 2002, **34**: 367-372.
49. Gonzales GF, Cordova A, Gonzales C, Chung A, Vega K, Villena A. ***Lepidium meyenii* (Maca) improved semen parameters in adult men.** *Asian J Androl* 2001, **3**: 301-303.
50. Obregon, L. *Maca*. Instituto de Fitoterapia Americana; 1998.
51. Bustos-Oregon E, Yucra S, Gonzales GF. ***Lepidium meyenii* (Maca) reduces spermatogenic damage induced by a single dose of malathion in mice.** *Asian J Androl* 2005, **7**: 71-76.
52. Farndale RW, Buttle DJ, Barrett AJ. **Improved quantitation and discrimination of sulphated glycosaminoglycans by use of dimethyl-methylene blue.** *Biochem Biophys Acta* 1986, **883**: 173-177.
53. Grounds MD. **Reasons for the degeneration of ageing skeletal muscle: a central role for IGF-1 signalling.** *Biogerontology* **3**: 19-24, 2002.
54. De Ceuninck F, Claez A, Dassencourt L, Anract P, Renard P. **Pharmacological disruption of insulin-like growth factor 1 binding to IGF-binding proteins restores anabolic responses in human osteoarthritic chondrocytes.** *Arthritis Res Ther* 2004, **6**: R393-R403.
55. Haupt JL, Frisbie DD, McIlwraith CM, Robbins PD, Ghivizzani S, Evans CH, Nixon A. **Dual transduction of insulin-like growth factor-I and interleukin-1 receptor antagonist protein controls cartilage degradation in an osteoarthritic culture model.** *J Orthopaedic Res* 2005, **23**: 118-126.

56. Iwanage H, Matsumoto T, Enomoto H, Okano K, Hishikawa Y, Shindo H, Koji T. **Enhanced expression of insulin-like growth factor-binding proteins in human osteoarthritic cartilage detected by immunohistochemistry and in situ hybridization.** *Osteoarthritis Cartilage* 2005, **13**: 439-448.
57. Verschure PJ, Van Noorden CJ, Van Marle J, Van de Berg WB. **Articular cartilage destruction in experimental inflammatory arthritis: insulin-like growth factor-1 regulation of proteoglycan metabolism in chondrocytes.** *Histochem J* 1998, **28**: 286-292.
58. Hui W, Rowan AD, Cawston TE. **Insulin-like growth factor 1 blocks collagen release and down regulates matrix metalloproteinase-1, -3, -8, and -13 mRNA expression in bovine nasal cartilage stimulated with oncostatin M in combination with interleukin1 $\alpha$ .** *Ann Rheum Dis* 2001, **60**: 254-261.
59. Rogachefsky RA, Dean DD, Howell DS, Altman RD. **Treatment of canine osteoarthritis with sodium pentosan polysulfate and insulin-like growth factor-1.** *Ann NY Acad Sci* 1994, **732**: 392-394.
60. Zofkova I. **Pathophysiological and clinical importance of insulin-like growth factor-1 with respect to bone metabolism.** *Physiol Rev* 2003, **52**: 657-679.
61. Chu CR, Kaplan LD, Fu FH, Crossett LS, Studer RK. **Recovery of articular cartilage metabolism following thermal stress is facilitated by IGF-1 and JNK inhibitor.** *Am J Sports Med* 2004, **32**: 191-196.
62. Tuncel M, Halici M, Canoz O, Turk CY, Oner M, Ozturk F, Kabak S. **Role of insulin like growth factor in repair response in immature cartilage.** *The Knee* 2005, **12**: 113-119.
63. Madry H, Kaul G, Cucchiari M, Stein U, Zurakowski D, Remberger K, Menger MD, Kohnn D, Trippel SB. **Enhanced repair of articular cartilage defects in**

- vivo by transplanted chondrocytes overexpressing insulin-like growth factor I (IGF-1).** *Gene Ther* 2005, **12**: 1171-1179.
64. Sandoval M, Okuhama NN, Angeles FM, Melchor VV, Condezo LA, Lao J, Miller MJS. **Antioxidant activity of the cruciferous vegetable maca (*Lepidium meyenii*).** *Food Chem* 2002, **79**: 207-213.
65. Studer RK. **Nitric oxide decreases IGF-1 receptor function in vitro: glutathione depletion enhances this effect in vivo.** *Osteoarthritis Cartilage* 2004, **12**: 863-869.

## Links

## Figure Legends

### **Figure 1: Enhanced expression of IGF-1 gene in human chondrocytes.**

IGF-1 gene expression referenced by the activity of  $\beta$ -actin was examined in human chondrocytes under basal (control) conditions and after treatment with RNI 249 50  $\mu\text{g/ml}$  (R50), vincaria 10  $\mu\text{g/ml}$  (V10) and their combination (R50 + V10). The \* refers to a significant difference from control levels ( $P < 0.05$ ), and the \*\* refers to a significant difference from all other groups ( $P < 0.001$ ), indicating that the combination of R50 and V10 produced significantly greater increases in IGF-1 expression than either V10 or R50 alone.

### **Figure 2: Production of IGF-1 from human cartilage explants as measured by media IGF-1 levels.**

RNI 249 produced a dose-dependent increase in media IGF-1 levels, as determined by ELISA. Only the 50 µg/ml concentration of RNI 249 produced a significant increase over basal, untreated explants (\* P<0.05, n=6).

**Figure 3: IGF-1 production from human cartilage explants in response to IL-1 $\beta$ , vincaria and RNI 249 and their combination.**


Control, untreated cartilage explants release a defined amount of IGF-1 into the bathing media. However, in IL-1 $\beta$  (5 ng/ml) treated explants IGF-1 levels are immeasurable (n=5). Co-treatment with either vincaria 10 µg/ml (V10, n=3), or RNI 249 50 µg/ml (R50, n=6), partially restored IGF-1 production. The combination of vincaria and RNI 249 (IL-1 + V +R, n=3) produced additive effects that resulted in the restoration of IGF-1 levels to control values despite the presence of IL-1 $\beta$ . The \* denotes a significant difference from all other groups (P<0.01).

**Figure 4: Normalization of IL-1 $\beta$  enhanced glycosaminoglycan release from human cartilage explants by RNI 249.**

Treatment of human cartilage explants (n=6) with IL-1 $\beta$  (5 ng/ml) results in the release of glycosaminoglycans (GAG) into the media. Administration of RNI 249 produced a dose-dependent decrease of IL-1 $\beta$  induced GAG release that was significant at the 10 µg/ml (R10, \* P<0.05) and 50 µg/ml (\*\* P<0.01) concentrations. The level of GAG release from control, untreated explants was indistinguishable from those explants treated with IL-1 $\beta$  + RNI 249 (50 µg/ml).

**Figure 5: Effects of vincaria and RNI 249 on basal and IL-1 $\beta$  stimulated nitrite production in human cartilage explants.**

Media nitrite levels, a reflection of nitric oxide production, released from human cartilage explants (n=3) was markedly enhanced by treatment with IL-1 $\beta$  (5 ng/ml, P<0.01). Co-treatment with either vincaria 10  $\mu$ g/ml (V10) or RNI 249 50  $\mu$ g/ml (R50) reduced nitrite levels although these changes were not significant. However, the combination of V10 and R50 produced additive effects and a significant reduction in nitrite levels (\* P<0.01 vs. IL-1 $\beta$ ). Both V10 and R50 alone, reduced basal nitrite levels when compared to control untreated values but these differences were not significant.



IGF-1 Gene Expression in Human Chondrocytes




Figure 2




Figure 3




Figure 4




Figure 5